

Cow's milk is beneficial for children as it provides essential nutrients for healthy bone and tooth development as well as growth.

Cow's milk should only be avoided if an allergy has been officially diagnosed by a recognised health professional. It is recommended that services request written documentation of cow's milk allergy diagnosis (from the health professional) before excluding dairy products.

If a child is allergic to cow's milk, alternative foods and drinks will be needed to meet the child's requirements for growth and development (such as calcium fortified soy drink).

01

Allergy action plan

Allergy action plans are recommended to advise staff what to do if a known allergen is ingested. The action plan should be developed with the child's family and treating team (doctor, allergist, paediatrician) and be approved and signed by a recognised health professional involved in their care.

On enrolment, centres should request written documentation of confirmed allergies from the child's treating team. Parents should document exactly what their child can and cannot tolerate to avoid confusion and this should be written on the allergy action plan.

CSpecialised anaphylaxis action plans are essential for children with anaphylactic reactions to cow's milk protein.

For information about developing an allergy action plan visit <u>Allergy policy and allergy action plans</u> or the Australasian Society of Clinical Immunology and Allergy (ASCIA)* website www.allergy.org.au





Early childhood services

02

Why avoid cow's milk

Parents may request their child avoid cow's milk for a variety of reasons. It is very important to determine the reason for avoidance because completely avoiding milk products may not be necessary unless a true allergy is diagnosed.

Dietary management of lactose intolerance is very different to the dietary management of cow's milk allergy (refer to the 'lactose intolerance' section of Food intolerance for more information). If a child is lactose intolerant but following the diet for a child with cow's milk allergy, they will be following an unnecessarily restrictive diet which can impact on growth and bone development. However, if the child has a true cow's milk allergy, this restrictive diet (with appropriate alternatives and supplementation) is necessary for their health.

Cow's milk is the most common cause of food allergies in children. Unless parents advise otherwise, children allergic to cow's milk should strictly avoid cow's milk and all foods containing cow's milk.



03

Avoiding cow's milk as a drink

If children are allergic to cow's milk, a calcium fortified alternative is required. The alternative chosen will depend on the child's age and whether or not the child tolerates soy drink. Approximately 40% of children who are allergic to cow's milk are also allergic to soy drink. The most suitable cow's milk substitute should be determined by the child's doctor or dietitian.

Cereal and nut milks such as rice and oat drinks should only be used if advised by the child's doctor or dietitian as these are low in energy, protein and other nutrients. In addition, not all varieties include added calcium.

Infant formula is usually based on cow's milk. Children under 1 year of age will require an alternative infant formula which should be offered under the supervision of a health professional.

The following table outlines milk alternatives and their suitability for different age groups.



Early childhood services

Milk or alternative	Suitability
Partially hydrolysed infant formula Examples: Nan HA Gold, Nan AR, Aptamil Gold + HA, Karicare Comfort	Not suitable for infants with established cow's milk allergy.
Soy infant formula Examples: Karicare Soy, S26 Gold Soy.	Recommended first alternative to milk for infants over 6 months of age Up to 40% of children who are allergic to cow's milk will also be allergic to soy products.
Extensively hydrolysed and amino acid based infant formula Examples: Aptamil Gold PeptiJunior, Alfare, Aptamil Gold Allerpro, Elecare, Neocate and Neocate Advance.	Suitable from birth to 2 years. Specialised hypoallergenic formula available only with prescription from medical practitioner.
Sheep and goats milk	Not suitable for infants under 12 months. Not a suitable alternative to cow's milk because these contain a similar protein to that found in cow's milk and can result in similar symptoms.
Soy drink	Not suitable as a drink for infants under 12 months. Suitable for older children (1-2 years) if soy is tolerated. Ensure calcium fortified soy products are used.
Cereal and nut milks Examples: Rice milk, oat milk, almond milk.	Not suitable as a drink for infants under 2 years of age. Usually well tolerated for children who react to cow's milk and soy drink. Low in fat and protein and need to be used with caution for children under 2 years of age or children with slow growth. Ensure calcium fortified varieties are used.
A2 milk	Not suitable for infants under 12 months. Not a suitable alternative to cow's milk because it contains cow's milk protein and can result in similar symptoms.
Lactose free milk	Not suitable for infants under 12 months. Not a suitable alternative to cow's milk because it contains cow's milk protein and can result in similar symptoms.

04

Avoiding cows milk in food

Avoidance of cow's milk and cow's milk products is required for children who have a cow's milk allergy. The following table outlines foods which are likely to contain cow's milk, and foods which are likely to be cow's milk free.



Early childhood services

Food group	Cow's milk free foods	Foods containing milk or likely to contain milk
Fruit and vegetables	Fresh, dried, canned fruit and vegetables Fruit and vegetable juices	Commercial potato salad, mashed potato, instant potato, fruit bars, vegetables in tins or packets, canned soups, canned vegetables in mayonnaise
Grains (breads and cereals)	Most bread Breakfast cereals (e.g. Rice Bubbles, Weetbix, Cornflakes) Plain baby rice cereal Wheat, oats, rice, bran, other grains Flours: wheat, corn, rice, soy, potato Pasta, spaghetti, noodles, sago, couscous, polenta, semolina, tapioca	Bread with milk products/cheese, commercial garlic bread French toast Glazed products Some muesli, muesli bars and chocolate flavoured cereals Commercial mixed baby cereals and muesli, baby rusks Canned spaghetti, macaroni cheese, filled pasta
Milk, yoghurt, cheese and alternatives and dairy desserts	Soy infant formula, soy drink (calcium fortified) Specialised infant formula Most soy yoghurt (check label) (may use milk based starter culture) Some soy cheeses (check label) Coconut based yoghurt and ice cream Custard and milk desserts made with soy or rice milk, soy ice confection	Milk: cow's, sheep, goat, lactose free, reduced fat, skim, UHT, A2, powdered, evaporated, condensed, buttermilk Regular and partially hydrolysed infant formula Yoghurt, Fruche, custard, yoghurt drinks, fermented milk drinks All cheese e.g. cheddar, ricotta, cream, cottage, goat Most vegetarian or soy cheeses (often have casein, a milk protein, as an ingredient) Sorbet, gelato, ice confection (check label)
Meat, poultry, fish, eggs, legumes and nuts	Fresh/frozen meat, chicken, fish Canned fish in oil, brine or water Beans/legumes (canned or dried), plain baked beans Tofu, eggs Nuts, nut paste, peanut butter	Some patties, burgers, pasties, sausages, meatloaf, fish fingers, chicken nuggets, sausage rolls, pies, processed meats, sandwich meats, crumbed/battered products (check label) Omelette, quiche, scrambled egg unless made with milk free products Flavoured baked beans (e.g. cheese in sauce)
Cakes and biscuits	Cakes, biscuits, muffins, slices, pancakes using milk free margarine and soy or rice drink	Commercial cakes, biscuits, slices, pastries unless labelled milk free Waffles, pancakes and pikelets Flavoured dry biscuits (check label)
Salad dressings and sauces	Vinegar based salad dressing e.g. French, Italian, balsamic Homemade/commercial gravies Tomato sauce, soy sauce	Creamy salad dressing e.g. coleslaw, Caesar, mayonnaise Béarnaise, tartare, hollandaise sauce Béchamel, white or cheese sauce
Soups	Homemade soup without milk products	Creamed soups
Jams/spreads	Vegemite, honey, jam, marmalade	Nutella, cheese spread
Drinks	Flavoured toppings, cocoa Soy or rice milk	Milk shakes, flavoured milk drinks, chocolate drink powders, malted milk powders, high protein drink powders, drinking chocolate
Fats and oils	Cooking oil, milk free margarine	Ghee, butterfat, margarine, butter, cream, imitation cream
Other	Tomato salsa, hummus dip Stock cubes, herbs/spices, salt Essences	Commercial dips (check label), yoghurt sauce, raita dip

Early childhood services



Avoiding cow's milk in packaged foods

In Australia, cow's milk must be declared on food labels. Labels should be carefully checked for presence of cow's milk (even if you have brought the product before) as product recipes can change.

The following ingredients indicate the presence of cow's milk in a product:

- milk (fresh, UHT, evaporated, condensed, powdered, A2, buttermilk), non fat milk solids, milk solids, milk powder
- butter, butter oil, butter fat, ghee
- whey, hydrolysed whey, casein, caseinate
- lactose
- chocolate

- beverage whitener
- lactalbumin
- cheese
- rennet, curds
- ice cream, cream
- · custard, yoghurt, fromage frais
- natural flavouring (cow's milk)



Replacing cow's milk in recipes

Try the simple substitutions below to make your recipes cow's milk free. Remember, many children who are allergic to cow's milk are also allergic to soy dripk

Check with the parents of allergic children before offering milk alternatives such as soy.



Ingredient	Allergy alternative
Cow's milk	Soy beverage or rice drink. Choose calcium fortified brands
Butter, margarine	Milk free margarine
Cheese	Omit
Crumbed meat or chicken	Grilled meat or chicken (uncrumbed)
Yoghurt	Soy yoghurt (highly allergic children may still react due to use of a milk based starter) Custard made with rice drink

^{*} The ASCIA website includes personal action plans for allergic reactions and for anaphylaxis. These are medical documents that can only be completed and signed by the patient's treating medical doctor and cannot be altered without their permission.

Adapted with permission from: Cow's Milk Allergy fact sheet, Department of Allergy and Immunology, Royal Children's Hospital Melbourne, April 2013.

Other references: Cow's milk free diet, Women's and Children's Health Network, Women's and Children's Hospital SA, March 2011.